

CORO7 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP55375**Specification**

CORO7 Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	P57737
Reactivity	Rat, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	100605

CORO7 Polyclonal Antibody - Additional Information**Gene ID** 79585**Other Names**

Coronin-7, Crn7, 70 kDa WD repeat tumor rejection antigen homolog, CORO7

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

CORO7 Polyclonal Antibody - Protein Information**Name** CORO7**Function**

F-actin regulator involved in anterograde Golgi to endosome transport: upon ubiquitination via 'Lys-33'-linked ubiquitin chains by the BCR(KLHL20) E3 ubiquitin ligase complex, interacts with EPS15 and localizes to the trans-Golgi network, where it promotes actin polymerization, thereby facilitating post-Golgi trafficking. May play a role in the maintenance of the Golgi apparatus morphology.

Cellular Location

Golgi apparatus membrane. Golgi apparatus, trans- Golgi network. Cytoplasmic vesicle. Cytoplasm, cytosol. Note=Predominantly cytosolic. Detected on vesicle-like cytoplasmic structures and on the cis-Golgi. Not associated with actin filaments

Tissue Location


Widely expressed. Expressed in the spleen, peripheral leukocytes, testes, brain, thymus and small intestine

CORO7 Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CORO7 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Human hepatocellular carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CORO7) Polyclonal Antibody, Unconjugated (bs-13999R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Human cervical cancer); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CORO7) Polyclonal Antibody, Unconjugated (bs-13999R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.