

COX18 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP55376**Specification**

COX18 Polyclonal Antibody - Product Information

| | |
|-------------------|------------------------|
| Application | IHC-P |
| Primary Accession | Q8N8Q8 |
| Reactivity | Rat, Pig, Dog, Bovine |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 37063 |

COX18 Polyclonal Antibody - Additional Information**Gene ID** 285521**Other Names**

Cytochrome c oxidase assembly protein COX18, mitochondrial, COX18Hs, Cytochrome c oxidase assembly protein 18, COX18, OXA1L2

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

COX18 Polyclonal Antibody - Protein Information**Name** COX18**Synonyms** OXA1L2**Function**

Mitochondrial membrane insertase required for the translocation of the C-terminus of cytochrome c oxidase subunit II (MT- CO2/COX2) across the mitochondrial inner membrane. Plays a role in MT-CO2/COX2 maturation following the COX20-mediated stabilization of newly synthesized MT-CO2/COX2 protein and before the action of the metallochaperones SCO1/2. Essential for the assembly and stability of the mitochondrial respiratory chain complex IV (also known as cytochrome c oxidase).

Cellular Location


Mitochondrion inner membrane; Multi-pass membrane protein

COX18 Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

COX18 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (COX18) Polyclonal Antibody, Unconjugated (bs-14006R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining..



Paraformaldehyde-fixed, paraffin embedded (Rat liver); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (COX18) Polyclonal Antibody, Unconjugated (bs-14006R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.