

CRISP10 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP55401**Specification**

CRISP10 Polyclonal Antibody - Product Information

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q9H336
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	56888

CRISP10 Polyclonal Antibody - Additional Information**Gene ID** 83690**Other Names**

Cysteine-rich secretory protein LCCL domain-containing 1, CocoaCrisp, Cysteine-rich secretory protein 10, CRISP-10, LCCL domain-containing cysteine-rich secretory protein 1, Trypsin inhibitor HI, CRISPLD1, CRISP10, LCRISP1

Dilution

WB~~1:1000<br \>IHC-P~~N/A<br \>IHC-F~~N/A<br \>IF~~1:50~200<br \>ICC~~N/A<br \>E~~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

CRISP10 Polyclonal Antibody - Protein Information**Name** CRISPLD1**Synonyms** CRISP10, LCRISP1**Cellular Location**


Secreted.

CRISP10 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CRISP10 Polyclonal Antibody - Images



Sample:


Heart(Mouse) Lysate at 40 ug

Primary: Anti-CRISP10 (bs-14060R) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution


Predicted band size: 54 kD

Observed band size: 54 kD



Paraformaldehyde-fixed, paraffin embedded (Human liver cancer); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CRISP10) Polyclonal Antibody, Unconjugated (bs-14060R) at 1:400 overnight at 4°C, followed by

operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Human stomach cancer); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CRISP10) Polyclonal Antibody, Unconjugated (bs-14060R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.