

MC2R Antibody (C-term)

Peptide Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP5540b

Specification

MC2R Antibody (C-term) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	Q01718
Other Accession	Q8HYN8 , Q64326 , NP_000520.1
Reactivity	Human
Predicted	Mouse, Pig
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	261-288

MC2R Antibody (C-term) - Additional Information

Gene ID 4158

Other Names

Adrenocorticotrophic hormone receptor, ACTH receptor, ACTH-R, Adrenocorticotropin receptor, Melanocortin receptor 2, MC2-R, MC2R, ACTHR

Target/Specificity

This MC2R antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 261-288 amino acids of human MC2R.

Dilution

WB~~1:1000
IHC-P~~1:10~50
FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.


Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.


Precautions

MC2R Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

MC2R Antibody (C-term) - Protein Information



MC2R Antibody (C-term) (Cat. #AP5540b) western blot analysis in WiDr cell line lysates (15ug/lane). This demonstrates the MC2R antibody detected the MC2R protein (arrow).



MC2R Antibody (C-term) (Cat. #AP5540b) immunohistochemistry analysis in formalin fixed and paraffin embedded human skin carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the MC2R Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

Name MC2R

Synonyms ACTHR

Function

Receptor for corticotropin (ACTH). This receptor is mediated by G proteins (G(s)) which activate adenylate cyclase (cAMP).

Cellular Location

Cell membrane; Multi-pass membrane protein.

Tissue Location

Melanocytes and corticoadrenal tissue.


MC2R Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MC2R Antibody (C-term) - Citations

- [Steroidogenic enzyme profile in an androgen-secreting adrenocortical oncocytoma associated with hirsutism.](#)



MC2R Antibody (C-term) (Cat. #AP5540b) flow cytometric analysis of WiDr cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

MC2R Antibody (C-term) - Background

MC2R encodes one member of the five-member G-protein associated melanocortin receptor family. Melanocortins (melanocyte-stimulating hormones and adrenocorticotrophic hormone) are peptides derived from pro-opiomelanocortin (POMC). MC2R is selectively activated by adrenocorticotrophic hormone, whereas the other four melanocortin receptors recognize a variety of melanocortin ligands.

MC2R Antibody (C-term) - References

Holliday, K.L., et al. Ann. Rheum. Dis. 69(3):556-560(2010)
Ding, Y.X., et al. Pharmacogenet. Genomics 20(2):71-76(2010)
Roy, S., et al. Endocrinology 151(2):660-670(2010)