

CSDE1 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP55415

Specification

CSDE1 Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	O75534
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	89 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human CSDE1
Epitope Specificity	501-600/798
Isotype	IgG
Purity	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm.
SIMILARITY	Contains 9 CSD (cold-shock) domains.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

UNR is a 798 amino acid protein that localizes to the cytoplasm and contains nine CDS (cold shock) domains. Existing as a component of the multi-protein autoregulatory ribonucleoprotein complex (ARC), UNR functions as an RNA-binding protein that is required for the initiation of rhinovirus RNA translation and is thought to be involved in translationally coupled mRNA turnover. UNR is expressed as two isoforms, designated long and short, and shares over 98% amino acid identity with its rat counterpart, suggesting a conserved role between species. The gene encoding UNR maps to human chromosome 1, which spans 260 million base pairs, contains over 3,000 genes and comprises nearly 8% of the human genome.

CSDE1 Polyclonal Antibody - Additional Information

Gene ID 7812

Other Names

Cold shock domain-containing protein E1, N-ras upstream gene protein, Protein UNR, CSDE1 (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=29905)
[HGNC:29905](#)

Dilution

IHC-P ~ N/A
IHC-F ~ N/A

="dilution_IF">IF~~1:50~200<br \>ICC~~N/A<br \>E~~N/A

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

CSDE1 Polyclonal Antibody - Protein Information

Name CSDE1 ([HGNC:29905](#))

Function

RNA-binding protein involved in translationally coupled mRNA turnover (PubMed:11051545, PubMed:15314026). Implicated with other RNA- binding proteins in the cytoplasmic deadenylation/translational and decay interplay of the FOS mRNA mediated by the major coding-region determinant of instability (mCRD) domain (PubMed:11051545, PubMed:15314026). Required for efficient formation of stress granules (PubMed:29395067).

Cellular Location


Cytoplasm. Cytoplasm, Stress granule. Cytoplasm, P-body

CSDE1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CSDE1 Polyclonal Antibody - Images



Tissue/cell: human lung carcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded;
Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-CSDE1 Polyclonal Antibody, Unconjugated(bs-14083R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: mouse embryo tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;
Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-CSDE1 Polyclonal Antibody, Unconjugated(bs-14083R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining