

DARS Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP55454

Specification

DARS Polyclonal Antibody - Product Information

Application WB, IHC-P, IHC-F, IF, ICC, E

Primary Accession P14868

Reactivity
Host
Rat, Pig, Dog, Bovine
Rabbit

Clonality Polyclonal
Calculated MW 57 KDa
Physical State Liquid

Immunogen KLH conjugated synthetic peptide derived

from human Cell proliferation inducing

protein 40 401-501/501

IgG

Epitope Specificity

Isotype Purity

affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Cytoplasm.

SIMILARITY Belongs to the class-II aminoacyl-tRNA

synthetase family.

Important Note This product as supplied is intended for

research use only, not for use in human,

therapeutic or diagnostic applications.

Background Descriptions

Aspartyl-tRNA synthetase (DARS) is part of a multienzyme complex of aminoacyl-tRNA synthetases. Aspartyl-tRNA synthetase charges its cognate tRNA with aspartate during protein biosynthesis. [provided by RefSeq, Jul 2008]

DARS Polyclonal Antibody - Additional Information

Gene ID 1615

Other Names

Aspartate--tRNA ligase, cytoplasmic, 6.1.1.12, Aspartyl-tRNA synthetase, AspRS, Cell proliferation-inducing gene 40 protein, DARS1 (HGNC:2678), DARS

Dilution

WB~~1:1000/>span class

="dilution IHC-P">IHC-P~~N/A<br \><span class

="dilution_IHC-F">IHC-F~~N/A<br \><span class

="dilution_IF">IF~~1:50~200<br \> < span class = "dilution_ICC">ICC~~N/A < br \> < span class = "dilution_ICC">ICC~~N/A < span class = "dilution_ICC">ICC~~N/A < span class = "dilution_ICC">ICC~



\>E~~N/A

Storage

Store at -20 $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 $^{\circ}$ C.

DARS Polyclonal Antibody - Protein Information

Name DARS1 (HGNC:2678)

Synonyms DARS

Function

Catalyzes the specific attachment of an amino acid to its cognate tRNA in a 2 step reaction: the amino acid (AA) is first activated by ATP to form AA-AMP and then transferred to the acceptor end of the tRNA.

Cellular Location

Cytoplasm, cytosol.

Tissue Location

Expression in the developing and adult brain shows similar patterns. Highly expressed in the ventricular and subventricular zones, including hippocampal subfields, the midlateral temporal cortex and the frontal polar cortex. The cerebellum, cerebral cortex, hippocampus, and lateral ventricle show preferential neuronal expression. Expression in the peripheral neurons is evident in the colon.


DARS Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

DARS Polyclonal Antibody - Images





Paraformaldehyde-fixed, paraffin embedded (Human liver cancer); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (DARS) Polyclonal Antibody, Unconjugated (bs-14197R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Sample:

Liver (Mouse) Lysate at 40 ug

Primary: Anti- DARS (bs-14197R) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 57 kD Observed band size: 57 kD