

**DAZAP2 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP55457****Specification****DAZAP2 Polyclonal Antibody - Product Information**

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	<a href="#">Q15038</a>
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	17319

**DAZAP2 Polyclonal Antibody - Additional Information****Gene ID** 9802**Other Names**

DAZ-associated protein 2, Deleted in azoospermia-associated protein 2, DAZAP2, KIAA0058

**Dilution**

dilution\_IHC-P">IHC-P~N/A  
dilution\_IHC-F">IHC-F~N/A  
dilution\_IF">IF~1:50~200  
dilution\_ICC">ICC~N/A  
dilution\_E">E~N/A

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**DAZAP2 Polyclonal Antibody - Protein Information****Name** DAZAP2 ([HGNC:2684](#))**Function**

In unstressed cells, promotes SIAH1-mediated polyubiquitination and degradation of the serine/threonine-protein kinase HIPK2, probably by acting as a loading factor that potentiates complex formation between HIPK2 and ubiquitin ligase SIAH1 (PubMed:[33591310](http://www.uniprot.org/citations/33591310)). In response to DNA damage, localizes to the nucleus following phosphorylation by HIPK2 and modulates the expression of a subset of TP53/p53 target genes by binding to TP53 at target gene promoters (PubMed:[33591310](http://www.uniprot.org/citations/33591310)). This limits the expression of a number of cell death-mediating TP53 target genes, reducing DNA damage-induced cell death (PubMed:[33591310](http://www.uniprot.org/citations/33591310)). Enhances the binding of transcription factor TCF7L2/TCF4, a Wnt signaling pathway effector, to the promoters of target genes (By similarity). Plays a role in stress granule formation (PubMed:[17984221](http://www.uniprot.org/citations/17984221)).

**Cellular Location**

Cytoplasm. Nucleus. Nucleus speckle. Nucleus, nuclear body. Cytoplasm, Stress granule  
Note=Predominantly nuclear in macrophages, stimulation of IL17RB with its ligand IL17E induces accumulation in the cytoplasm (PubMed:22070932). Predominantly cytoplasmic when unphosphorylated and localizes to the nucleus following phosphorylation by HIPK2 (PubMed:33591310). Localizes to stress granules under cellular stress conditions (PubMed:17984221).

**Tissue Location**

Widely expressed. Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon and leukocytes. Down- regulated in multiple myeloma.

**DAZAP2 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**DAZAP2 Polyclonal Antibody - Images**