

**DDHD1 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP55468****Specification**

---

**DDHD1 Polyclonal Antibody - Product Information**

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	<a href="#">Q8NEL9</a>
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	100 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human DDHD1
Epitope Specificity	751-850/900
Isotype	IgG
<b>Purity</b>	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasmic
SIMILARITY	Belongs to the PA-PLA1 family.Contains 1 DDHD domain.
SUBUNIT	Forms homooligomers and, to a much smaller extent, heterooligomers with DDHD2.
DISEASE	Spastic paraplegia 28, autosomal recessive (SPG28) [MIM:609340]: A form of spastic paraplegia, a neurodegenerative disorder characterized by a slow, gradual, progressive weakness and spasticity of the lower limbs. Rate of progression and the severity of symptoms are quite variable. Initial symptoms may include difficulty with balance, weakness and stiffness in the legs, muscle spasms, and dragging the toes when walking. In some forms of the disorder, bladder symptoms (such as incontinence) may appear, or the weakness and stiffness may spread to other parts of the body. Some SPG28 patients also have distal sensory impairment. Note=The disease is caused by mutations affecting the gene represented in this entry.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

**Background Descriptions**

Phosphatidic acid is released following cell activation and functions as a second messenger in several signaling pathways. DDHD1 is a lipase that catalyzes degradation of phosphatidic acid and attenuates cell activation.

## DDHD1 Polyclonal Antibody - Additional Information

**Gene ID** 80821

### Other Names

Phospholipase DDHD1, 3.1.1.-, DDHD domain-containing protein 1, Phosphatidic acid-preferring phospholipase A1 homolog, PA-PLA1, DDHD1 ([http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?hgnc\\_id=19714](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=19714)), KIAA1705

### Target/Specificity

Highly expressed in testis. Also expressed in brain, spleen and lung. Only expressed in cerebellum in fetal brain.

### Dilution

IHC-P ~ N/A  
IHC-F ~ N/A  
IF ~ 1:50 ~ 200  
ICC ~ N/A  
E ~ N/A

### Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

### Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

## DDHD1 Polyclonal Antibody - Protein Information

**Name** DDHD1 ([HGNC:19714](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=19714))

**Synonyms** KIAA1705

### Function

Phospholipase A1 (PLA1) that hydrolyzes ester bonds at the sn-1 position of glycerophospholipids producing a free fatty acid and a lysophospholipid (Probable) (PubMed: [20359546](http://www.uniprot.org/citations/20359546), PubMed: [22922100](http://www.uniprot.org/citations/22922100)). Prefers phosphatidate (1,2-diacyl-sn-glycero-3-phosphate, PA) as substrate in vitro, but can efficiently hydrolyze phosphatidylinositol (1,2-diacyl-sn-glycero-3-phospho-(1D-myo-inositol), PI), as well as a range of other glycerophospholipid substrates such as phosphatidylcholine (1,2-diacyl-sn-glycero-3-phosphocholine, PC), phosphatidylethanolamine (1,2-diacyl-sn-glycero-3-phosphoethanolamine, PE), phosphatidylserine (1,2-diacyl-sn-glycero-3-phospho-L-serine, PS) and phosphatidylglycerol (1,2-diacyl-sn-glycero-3-phospho-(1'-sn-glycerol), PG) (Probable) (PubMed: [20359546](http://www.uniprot.org/citations/20359546)). Involved in the regulation of the endogenous content of polyunsaturated PI and PS lipids in the nervous system. Changes in these lipids extend to downstream metabolic products like PI phosphates PIP and PIP2, which play fundamental roles in cell biology (By similarity). Regulates mitochondrial

morphology (PubMed:<a href="http://www.uniprot.org/citations/24599962" target="\_blank">24599962</a>). These dynamic changes may be due to PA hydrolysis at the mitochondrial surface (PubMed:<a href="http://www.uniprot.org/citations/24599962" target="\_blank">24599962</a>). May play a regulatory role in spermatogenesis or sperm function (PubMed:<a href="http://www.uniprot.org/citations/24599962" target="\_blank">24599962</a>).

**Cellular Location**

Cytoplasm.

**Tissue Location**

Highly expressed in testis. Also expressed in brain, spleen and lung. Only expressed in cerebellum in fetal brain

**DDHD1 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**DDHD1 Polyclonal Antibody - Images**