

DENND4A Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP55496**Specification**

DENND4A Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF, ICC
Primary Accession	Q7Z401
Reactivity	Rat, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	209244

DENND4A Polyclonal Antibody - Additional Information**Gene ID** 10260**Other Names**

C-myc promoter-binding protein, DENN domain-containing protein 4A, DENND4A, IRLB, MYCPBP

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

DENND4A Polyclonal Antibody - Protein Information**Name** DENND4A**Synonyms** IRLB, MYCPBP**Function**

Probable guanine nucleotide exchange factor (GEF) which may activate RAB10. Promotes the exchange of GDP to GTP, converting inactive GDP-bound Rab proteins into their active GTP-bound form. According to PubMed:8056341, it may bind to ISRE-like element (interferon-stimulated response element) of MYC P2 promoter.

Cellular Location

Nucleus.

Tissue Location

Expressed ubiquitously. Highest expression in bone marrow, medium in peripheral blood lymphocytes and lowest in spleen. In brain, breast, and prostate, higher expression was seen in normal cells than in tumor cells. Expression is regulated in a growth- and cell cycle-dependent manner.

DENND4A Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DENND4A Polyclonal Antibody - Images