

Dermatopontin Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP55504**Specification**

Dermatopontin Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q07507
Reactivity	Rat, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	24 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human Dermatopontin
Epitope Specificity	101-201/201
Isotype	IgG
Purity	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Secreted > extracellular space > extracellular matrix.
SIMILARITY	Belongs to the dermatopontin family.
Post-translational modifications	Sulfated on tyrosine residue(s).
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

Dermatopontin is an extracellular matrix protein with possible functions in cell-matrix interactions and matrix assembly. The protein is found in various tissues and many of its tyrosine residues are sulphated. Dermatopontin is postulated to modify the behavior of TGF-beta through interaction with decorin. [provided by RefSeq, Jul 2008]

Dermatopontin Polyclonal Antibody - Additional Information**Gene ID** 1805**Other Names**

Dermatopontin, Tyrosine-rich acidic matrix protein, TRAMP, DPT

Target/Specificity

Expressed in fibroblasts, heart, skeletal muscle, brain and pancreas. Expressed at an intermediate level in lung and kidney, and at a low level in liver and placenta. Expressed at a lower level in fibroblasts from hypertrophic scar lesional skin and in fibroblasts from patients with systemic sclerosis than in normal skin fibroblasts.

Dilution

IHC-P~~N/A<br \>IHC-F~~N/A<br \>IF~~1:50~200<br \>ICC~~N/A<br \>E~~N/A

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Dermatopontin Polyclonal Antibody - Protein Information**Name** DPT**Function**

Seems to mediate adhesion by cell surface integrin binding. May serve as a communication link between the dermal fibroblast cell surface and its extracellular matrix environment. Enhances TGFβ1 activity. Inhibits cell proliferation. Accelerates collagen fibril formation, and stabilizes collagen fibrils against low-temperature dissociation (By similarity).

Cellular Location

Secreted, extracellular space, extracellular matrix

Tissue Location

Expressed in fibroblasts, heart, skeletal muscle, brain and pancreas. Expressed at an intermediate level in lung and kidney, and at a low level in liver and placenta. Expressed at a lower level in fibroblasts from hypertrophic scar lesional skin and in fibroblasts from patients with systemic sclerosis than in normal skin fibroblasts.

Dermatopontin Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Dermatopontin Polyclonal Antibody - Images