

DGKG Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP55510**Specification**

DGKG Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	P49619
Reactivity	Rat, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	89124

DGKG Polyclonal Antibody - Additional Information**Gene ID** 1608**Other Names**

Diacylglycerol kinase gamma, DAG kinase gamma, 2.7.1.107, Diglyceride kinase gamma, DGK-gamma, DGKG, DAGK3

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

DGKG Polyclonal Antibody - Protein Information**Name** DGKG**Synonyms** DAGK3**Function**

Diacylglycerol kinase that converts diacylglycerol/DAG into phosphatidic acid/phosphatidate/PA and regulates the respective levels of these two bioactive lipids (PubMed:8034597). Thereby, acts as a central switch between the signaling pathways activated by these second messengers with different cellular targets and opposite effects in numerous biological processes (By similarity). Has no apparent specificity with regard to the acyl compositions of diacylglycerol (PubMed:8034597). Specifically expressed in the cerebellum where it controls the level of diacylglycerol which in turn regulates the activity of protein kinase C gamma. Through protein kinase C gamma, indirectly regulates the dendritic development of Purkinje cells, cerebellar long term depression and ultimately cerebellar motor coordination (By similarity).

Cellular Location

Membrane. Cytoplasm, cytosol. Cytoplasm, cytoskeleton {ECO:0000250|UniProtKB:P49620}

Tissue Location


Predominantly expressed in retina and in a much lesser extent in the brain. Other tissues contain extremely low levels of DGK-gamma

DGKG Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DGKG Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (DGKG) Polyclonal Antibody, Unconjugated (bs-14297R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.