

DHI1L Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP55513**Specification**

DHI1L Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	Q7Z5J1
Host	Rabbit
Clonality	Polyclonal
Calculated MW	34288

DHI1L Polyclonal Antibody - Additional Information**Gene ID** 374875**Other Names**

Hydroxysteroid 11-beta-dehydrogenase 1-like protein, 1.1.1.-, 11-beta-hydroxysteroid dehydrogenase type 3, 11-DH3, 11-beta-HSD3, Short chain dehydrogenase/reductase family 26C member 2, Short-chain dehydrogenase/reductase 10, HSD11B1L, HSD3, SCDR10, SDR26C2

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

DHI1L Polyclonal Antibody - Protein Information**Name** HSD11B1L**Synonyms** HSD3, SCDR10, SDR26C2**Cellular Location**

Secreted.


DHI1L Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)

- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DHI1L Polyclonal Antibody - Images



Tissue/cell: human kidney tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;
Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;
Incubation: Anti-DHI1L Polyclonal Antibody, Unconjugated(bs-14303R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Paraformaldehyde-fixed, paraffin embedded (Human brain glioma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (DHI1L) Polyclonal Antibody, Unconjugated (bs-14303R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.