

**DHX37 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP55523****Specification****DHX37 Polyclonal Antibody - Product Information**

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	<a href="#">Q8IY37</a>
Reactivity	Rat, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	129545

**DHX37 Polyclonal Antibody - Additional Information****Gene ID** 57647**Other Names**

Probable ATP-dependent RNA helicase DHX37, 3.6.4.13, DEAH box protein 37, DHX37, DDX37, KIAA1517

**Dilution**

WB~1:1000  
IHC-P~N/A  
IHC-F~N/A  
IF~1:50~200  
ICC~N/A  
E~N/A

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**DHX37 Polyclonal Antibody - Protein Information****Name** DHX37 ([HGNC:17210](#))**Synonyms** DDX37, KIAA1517**Function**

ATP-binding RNA helicase that plays a role in maturation of the small ribosomal subunit in ribosome biogenesis (PubMed:<a href="http://www.uniprot.org/citations/30582406" target="\_blank">30582406</a>). Required for the release of the U3 snoRNP from pre-ribosomal particles (PubMed:<a href="http://www.uniprot.org/citations/30582406" target="\_blank">30582406</a>). Part of the small subunit (SSU) processome, first precursor of the small eukaryotic ribosomal subunit. During the assembly of the SSU processome in the

nucleolus, many ribosome biogenesis factors, an RNA chaperone and ribosomal proteins associate with the nascent pre-rRNA and work in concert to generate RNA folding, modifications, rearrangements and cleavage as well as targeted degradation of pre-ribosomal RNA by the RNA exosome (PubMed:<a href="http://www.uniprot.org/citations/34516797" target="\_blank">34516797</a>). Plays a role in early testis development (PubMed:<a href="http://www.uniprot.org/citations/31287541" target="\_blank">31287541</a>, PubMed:<a href="http://www.uniprot.org/citations/31337883" target="\_blank">31337883</a>). Probably also plays a role in brain development (PubMed:<a href="http://www.uniprot.org/citations/31256877" target="\_blank">31256877</a>).

**Cellular Location**

Nucleus, nucleolus. Cytoplasm. Nucleus membrane


**Tissue Location**

Expressed in the fallopian tube, ovary, uterus and testis. Also expressed in the brain.


**DHX37 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**DHX37 Polyclonal Antibody - Images**

Paraformaldehyde-fixed, paraffin embedded (rat brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (DHX37) Polyclonal Antibody, Unconjugated (bs-14320R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Sample: 293T Cell (Human) Lysate at 40 ug  
Primary: Anti-DHX37(bs-14320R) at 1/300 dilution  
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution  
Predicted band size: 129kD  
Observed band size: 135kD