

DPRX Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP55567**Specification****DPRX Polyclonal Antibody - Product Information**

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	A6NEQ7
Host	Rabbit
Clonality	Polyclonal
Calculated MW	22 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human DPRX
Epitope Specificity	1-100/191
Isotype	IgG
Purity	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nucleus.
SIMILARITY	Belongs to the paired homeobox family. Contains 1 homeobox DNA-binding domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

Homeobox genes encode DNA-binding proteins, many of which are thought to be involved in early embryonic development. Homeobox genes encode a DNA-binding domain of 60 to 63 amino acids referred to as the homeodomain. This gene is a member of the DPRX homeobox gene family. Evidence of mRNA expression has not yet been found for this gene. Multiple, related processed pseudogenes have been found which are thought to reflect expression of this gene in the germ line or embryonic cells. [provided by RefSeq, Jul 2008]

DPRX Polyclonal Antibody - Additional Information**Gene ID** 503834**Other Names**

Divergent paired-related homeobox, DPRX

Dilution

WB~~1:1000<br \>IHC-P~~N/A<br \>IHC-F~~N/A<br \>IF~~1:50~200<br \>ICC~~N/A<br \>E~~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

DPRX Polyclonal Antibody - Protein Information

Name DPRX ([HGNC:32166](#))

Function

Transcription factor that acts as a repressor.

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00108}.

DPRX Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DPRX Polyclonal Antibody - Images