

**EDF1 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP55604****Specification**

---

**EDF1 Polyclonal Antibody - Product Information**

Application	IHC-P, IHC-F, IF, ICC
Primary Accession	<a href="#">O60869</a>
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	16369

**EDF1 Polyclonal Antibody - Additional Information****Gene ID** 8721**Other Names**

Endothelial differentiation-related factor 1, EDF-1, Multiprotein-bridging factor 1, MBF1, EDF1

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**EDF1 Polyclonal Antibody - Protein Information****Name** EDF1**Function**

Transcriptional coactivator stimulating NR5A1 and ligand- dependent NR1H3/LXRA and PPARG transcriptional activities. Enhances the DNA-binding activity of ATF1, ATF2, CREB1 and NR5A1. Regulates nitric oxid synthase activity probably by sequestering calmodulin in the cytoplasm. May function in endothelial cells differentiation, hormone- induced cardiomyocytes hypertrophy and lipid metabolism.

**Cellular Location**

Cytoplasm. Nucleus. Note=Also nuclear upon binding to NR5A1 and treatment of cells with TPA or forskolin

**Tissue Location**

Expressed in brain, liver, lung, kidney and heart (at protein level). Ubiquitously expressed. More abundant in heart, pancreas, liver, intestine and adipose tissues

**EDF1 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**EDF1 Polyclonal Antibody - Images**