

EFR3B Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP55608**Specification****EFR3B Polyclonal Antibody - Product Information**

Application	IHC-P, IHC-F, IF, ICC
Primary Accession	O9Y2G0
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	92 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human EFR3B
Epitope Specificity	21-120/817
Isotype	IgG
Purity	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the EFR3 family.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

EFR3B (EFR3 homolog B) is an 817 amino acid protein that exists as three alternatively spliced isoforms and belongs to the EFR3 family. The gene encoding EFR3B maps to human chromosome 2p23.3 and mouse chromosome 12 A1.1. Human chromosome 2 is the second largest human chromosome, which consists of 237 million bases, encodes over 1,400 genes and makes up approximately 8% of the human genome. A number of genetic diseases are linked to genes on chromosome 2. Harlequin ichthyosis, a rare and morbid skin deformity, is associated with mutations in the ABCA12 gene. The lipid metabolic disorder sitosterolemia is associated with ABCG5 and ABCG8. An extremely rare recessive genetic disorder, Alström syndrome, is due to mutations in the ALMS1 gene. Interestingly, chromosome 2 contains what appears to be a vestigial second centromere and vestigial telomeres which gives credence to the hypothesis that human chromosome 2 is the result of an ancient fusion of two ancestral chromosomes seen in modern form today in apes.

EFR3B Polyclonal Antibody - Additional Information**Gene ID 22979****Other Names**

Protein EFR3 homolog B, EFR3B (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=29155)
HGNC:29155

Dilution

IHC-P ~ ~ N/A
 IHC-F ~ ~ N/A
 IF ~ ~ 1:50 ~ 200
 ICC ~ ~ N/A

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

EFR3B Polyclonal Antibody - Protein Information

Name EFR3B ([HGNC:29155](#))

Function

Component of a complex required to localize phosphatidylinositol 4-kinase (PI4K) to the plasma membrane (PubMed:[23229899](http://www.uniprot.org/citations/23229899) target="_blank">23229899, PubMed:[25608530](http://www.uniprot.org/citations/25608530) target="_blank">25608530, PubMed:[26571211](http://www.uniprot.org/citations/26571211) target="_blank">26571211). The complex acts as a regulator of phosphatidylinositol 4-phosphate (PtdIns(4)P) synthesis (Probable). In the complex, EFR3B probably acts as the membrane-anchoring component (PubMed:[23229899](http://www.uniprot.org/citations/23229899) target="_blank">23229899). Also involved in responsiveness to G-protein-coupled receptors; it is however unclear whether this role is direct or indirect (PubMed:[25380825](http://www.uniprot.org/citations/25380825) target="_blank">25380825).


Cellular Location


Cell membrane; Lipid-anchor. Cytoplasm, cytosol. Note=Palmitoylation anchors the protein to the plasma membrane (PubMed:23229899, PubMed:25380825). A small amount is observed in the cytosol (PubMed:25380825)

EFR3B Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

EFR3B Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (rat brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (EFR3B) Polyclonal Antibody, Unconjugated (bs-14521R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (EFR3B) Polyclonal Antibody, Unconjugated (bs-14521R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.