

EIF2D Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP55616

Specification

EIF2D Polyclonal Antibody - Product Information

Application Primary Accession Reactivity Host Clonality Calculated MW IHC-P, IHC-F, IF, ICC, E <u>P41214</u> Rat, Pig, Bovine Rabbit Polyclonal 64706

EIF2D Polyclonal Antibody - Additional Information

Gene ID 1939

Other Names Eukaryotic translation initiation factor 2D, eIF2d, Hepatocellular carcinoma-associated antigen 56, Ligatin, EIF2D, HCA56, LGTN

Dilution IHC-P~~N/A<br \>IHC-F~~N/A<br \>IF~~1:50~200<br \>ICC~~N/A<br \>E~~N/A

Format 0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

EIF2D Polyclonal Antibody - Protein Information

Name EIF2D

Synonyms HCA56, LGTN

Function

Translation initiation factor that is able to deliver tRNA to the P-site of the eukaryotic ribosome in a GTP-independent manner. The binding of Met-tRNA(I) occurs after the AUG codon finds its position in the P-site of 40S ribosomes, the situation that takes place during initiation complex formation on some specific RNAs. Its activity in tRNA binding with 40S subunits does not require the presence of the aminoacyl moiety. Possesses the unique ability to deliver non-Met (elongator) tRNAs into the P-site of the 40S subunit. In addition to its role in initiation, can promote release of deacylated tRNA and mRNA from recycled 40S subunits following ABCE1-mediated dissociation of



post-termination ribosomal complexes into subunits.

Cellular Location Cytoplasm.

EIF2D Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- <u>Flow Cytomety</u>
- <u>Cell Culture</u>

EIF2D Polyclonal Antibody - Images