

CUG-BP1/2 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP55623**Specification****CUG-BP1/2 Polyclonal Antibody - Product Information**

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	O95319
Reactivity	Rat, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	54285

CUG-BP1/2 Polyclonal Antibody - Additional Information**Gene ID** 10659**Other Names**

CUGBP Elav-like family member 2, CELF-2, Bruno-like protein 3, CUG triplet repeat RNA-binding protein 2, CUG-BP2, CUG-BP- and ETR-3-like factor 2, ELAV-type RNA-binding protein 3, ETR-3, Neuroblastoma apoptosis-related RNA-binding protein, hNAPOR, RNA-binding protein BRUNOL-3, CELF2, BRUNOL3, CUGBP2, ETR3, NAPOR

Dilution

WB~~1:1000<br \>IHC-P~~N/A<br \>IHC-F~~N/A<br \>IF~~1:50~200<br \>ICC~~N/A<br \>E~~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

CUG-BP1/2 Polyclonal Antibody - Protein Information**Name** CELF2**Synonyms** BRUNOL3, CUGBP2, ETR3, NAPOR**Function**

RNA-binding protein implicated in the regulation of several post-transcriptional events. Involved in pre-mRNA alternative splicing, mRNA translation and stability. Mediates exon inclusion and/or exclusion in pre-mRNA that are subject to tissue-specific and developmentally regulated alternative splicing. Specifically activates exon 5 inclusion of TNNT2 in embryonic, but not adult,

skeletal muscle. Activates TNNT2 exon 5 inclusion by antagonizing the repressive effect of PTB. Acts both as an activator and as a repressor of a pair of coregulated exons: promotes inclusion of the smooth muscle (SM) exon but exclusion of the non-muscle (NM) exon in actinin pre-mRNAs. Promotes inclusion of exon 21 and exclusion of exon 5 of the NMDA receptor R1 pre-mRNA. Involved in the apoB RNA editing activity. Increases COX2 mRNA stability and inhibits COX2 mRNA translation in epithelial cells after radiation injury (By similarity). Modulates the cellular apoptosis program by regulating COX2-mediated prostaglandin E2 (PGE2) expression (By similarity). Binds to (CUG)_n triplet repeats in the 3'-UTR of transcripts such as DMPK. Binds to the muscle-specific splicing enhancer (MSE) intronic sites flanking the TNNT2 alternative exon 5. Binds preferentially to UG-rich sequences, in particular UG repeat and UGUU motifs. Binds to apoB mRNA, specifically to AU-rich sequences located immediately upstream of the edited cytidine. Binds AU-rich sequences in the 3'-UTR of COX2 mRNA (By similarity). Binds to an intronic RNA element responsible for the silencing of exon 21 splicing (By similarity). Binds to (CUG)_n repeats (By similarity). May be a specific regulator of miRNA biogenesis. Binds to primary microRNA pri-MIR140 and, with CELF1, negatively regulates the processing to mature miRNA (PubMed:28431233).

Cellular Location

Nucleus. Cytoplasm {ECO:0000250|UniProtKB:Q7T2T1, ECO:0000250|UniProtKB:Q9Z0H4}
Note=Accumulates in the cytoplasm after ionizing radiation (By similarity). Colocalizes with APOBEC1 and A1CF. RNA-binding activity is detected in both nuclear and cytoplasmic compartments.

Tissue Location

Expressed in frontal cortex. Isoform 1 is expressed in brain and lung. Isoform 2 is expressed in heart, brain, placenta, lung, liver, kidney, skeletal muscle and pancreas. Isoform 4 is expressed in heart, lung, skeletal muscle, kidney and pancreas


CUG-BP1/2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CUG-BP1/2 Polyclonal Antibody - Images






Tissue/cell: human lung carcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-CUG-BP1/2 Polyclonal Antibody, Unconjugated(bs-14562R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining




Protein: spleen(mouse) lysate at 40ug;

Primary: rabbit Anti-CUG-BP1/2 (bs-14562R) at 1:300;


Secondary: HRP conjugated Goat-Anti-rabbit IgG(bs-0295G-HRP) at 1: 5000;

Predicted band size: 54 kD


Observed band size: 54 kD



Paraformaldehyde-fixed, paraffin embedded (rat lung); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CUG-BP1,2) Polyclonal Antibody, Unconjugated (bs-14562R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CUG-BP1,2) Polyclonal Antibody, Unconjugated (bs-14562R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat heart); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CUG-BP1,2) Polyclonal Antibody, Unconjugated (bs-14562R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.