

Emil Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP55632

Specification

Emi1 Polyclonal Antibody - Product Information

Application IHC-P, IHC-F, IF, ICC, E

Primary Accession <u>Q9UKT4</u>

Reactivity
Host
Clonality
Calculated MW
Rat, Pig, Dog, Bovine
Rabbit
Polyclonal
50 KDa

Calculated MW 50 KDa
Physical State Liquid

Immunogen KLH conjugated synthetic peptide derived

laG

from human Emi1

Epitope Specificity 101-200/447

Isotype
Purity
affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

SUBCELLULAR LOCATION

Proclin300 and 50% Glycerol.

Nucleus. Cytoplasm. Cytoplasm >
cytoskeleton > spindle. In interphase,
localizes in a punctate manner in the

nucleus and cytoplasm with some perinuclear concentration. In mitotic cells, localizes throughout the cell, particularly

at the spindle.

SIMILARITY Contains 1 F-box domain. Contains 1

IBR-type zinc finger.

Post-translational modifications Phosphorylation by CDK2 and

subsequently by PLK1 triggers degradation

during early mitosis through

ubiquitin-mediated proteolysis by the SCF ubiquitin ligase complex containing the F-box protein BTRC. This degradation is necessary for the activation of APC in late

mitosis and subsequent mitotic

progression.

Important Note

This product as supplied is intended for research use only, not for use in human,

therapeutic or diagnostic applications.

Background Descriptions

This gene encodes a member of the F-box protein family which is characterized by an approximately 40 amino acid motif, the F-box. The F-box proteins constitute one of the four subunits of the ubiquitin protein ligase complex called SCFs (SKP1-cullin-F-box), which function in phosphorylation-dependent ubiquitination. The F-box proteins are divided into 3 classes: Fbws containing WD-40 domains, Fbls containing leucine-rich repeats, and Fbxs containing either different protein-protein interaction modules or no recognizable motifs. The protein encoded by this gene belongs to the Fbxs class. This protein is similar to xenopus early mitotic inhibitor-1



(Emi1), which is a mitotic regulator that interacts with Cdc20 and inhibits the anaphase promoting complex. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Dec 2008]

Emi1 Polyclonal Antibody - Additional Information

Gene ID 26271

Other Names

F-box only protein 5, Early mitotic inhibitor 1, FBXO5 (HGNC:13584)

Dilution

```
<span class ="dilution_IHC-P">IHC-P~~N/A</span><br \> <span class
="dilution_IHC-F">IHC-F~~N/A</span><br \> <span class
="dilution_IF">IF~~1:50~200</span><br \> <span class ="dilution_ICC">ICC~~N/A</span><br \> <span class = "dilution_E">E~~N/A</span>
```

Storage

Store at -20 $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 $^{\circ}$ C.

Emi1 Polyclonal Antibody - Protein Information

Name FBXO5 (HGNC:13584)

Function

Regulator of APC activity during mitotic and meiotic cell cycle (PubMed:16921029, PubMed:17234884, PubMed:17485488, PubMed:17875940, PubMed:23708001, PubMed:23708605). During mitotic cell cycle plays a role as both substrate and inhibitor of APC-FZR1 complex (PubMed: 16921029, PubMed:17234884, PubMed:17485488, PubMed:17875940, PubMed:23708001, PubMed:23708605, PubMed:29875408). During G1 phase, plays a role as substrate of APC-FZR1 complex E3 ligase (PubMed:29875408). Then switches as an inhibitor of APC-FZR1 complex during S and G2 leading to cell-cycle commitment (PubMed:29875408). As APC inhibitor, prevents the degradation of APC substrates at multiple levels: by interacting with APC and blocking access of APC substrates to the D-box coreceptor, formed by FZR1 and ANAPC10; by suppressing ubiquitin ligation and chain elongation by APC by preventing the UBE2C and UBE2S activities (PubMed:16921029, PubMed:23708001, PubMed:<a href="http://www.uniprot.org/citations/23708605"



target="_blank">23708605). Plays a role in genome integrity preservation by coordinating DNA replication with mitosis through APC inhibition in interphase to stabilize CCNA2 and GMNN in order to promote mitosis and prevent rereplication and DNA damage-induced cellular senescence (PubMed:17234884, PubMed:17485488, PubMed:17875940). During oocyte maturation, plays a role in meiosis through inactivation of APC-FZR1 complex. Inhibits APC through RPS6KA2 interaction that increases FBXO5 affiniy for CDC20 leading to the metaphase arrest of the second meiotic division before fertilization (By similarity). Controls entry into the first meiotic division through inactivation of APC-FZR1 complex (By similarity). Promotes migration and osteogenic differentiation of mesenchymal stem cells (PubMed:29850565).

Cellular Location

Nucleus. Cytoplasm. Cytoplasm, cytoskeleton, spindle. Note=In interphase, localizes in a punctate manner in the nucleus and cytoplasm with some perinuclear concentration (PubMed:11988738). In mitotic cells, localizes throughout the cell, particularly at the spindle (PubMed:15469984)

Emi1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Emi1 Polyclonal Antibody - Images