

FAM108A1 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP55676

Specification

FAM108A1 Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF, ICC
Primary Accession	Q96GS6
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	32 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human FAM108A1
Epitope Specificity	251-310/310
Isotype	IgG
Purity	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Secreted
SIMILARITY	Belongs to the AB hydrolase superfamily. ABHD17 family.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

Belongs to the AB hydrolase superfamily. FAM108 family.

FAM108A1 Polyclonal Antibody - Additional Information

Gene ID 81926

Other Names

Alpha/beta hydrolase domain-containing protein 17A, Abhydrolase domain-containing protein 17A {ECO:0000312|HGNC:HGNC:28756}, 3.1.2.22, ABHD17A (HGNC:28756), C19orf27, FAM108A1

Dilution

IHC-P~~N/A<br \>IHC-F~~N/A<br \>IF~~1:50~200<br \>ICC~~N/A

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

FAM108A1 Polyclonal Antibody - Protein Information

Name ABHD17A ([HGNC:28756](#))

Synonyms C19orf27, FAM108A1

Function

Hydrolyzes fatty acids from S-acylated cysteine residues in proteins (PubMed:26701913). Has depalmitoylating activity towards NRAS (PubMed:26701913). Has depalmitoylating activity towards DLG4/PSD95 (PubMed:26701913). May have depalmitoylating activity towards MAP6 (By similarity).

Cellular Location


Cell membrane; Lipid-anchor; Cytoplasmic side. Endosome membrane; Lipid-anchor; Cytoplasmic side. Cell projection, dendritic spine {ECO:0000250|UniProtKB:Q5XIJ5}. Postsynaptic density membrane {ECO:0000250|UniProtKB:Q5XIJ5}

FAM108A1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FAM108A1 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Rat stomach); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (FAM108A1) Polyclonal Antibody, Unconjugated (bs-14691R) at 1:500 overnight at 4°C, followed

by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (FAM108A1) Polyclonal Antibody, Unconjugated (bs-14691R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.