

**FAM193B Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP55713****Specification**

---

**FAM193B Polyclonal Antibody - Product Information**

Application	IHC-P
Primary Accession	<a href="#">Q96PV7</a>
Reactivity	Rat, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	96543

**FAM193B Polyclonal Antibody - Additional Information****Other Names**

Protein FAM193B, FAM193B, IRIZIO, KIAA1931

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**FAM193B Polyclonal Antibody - Protein Information**

**Name** FAM193B

**Synonyms** IRIZIO, KIAA1931

**Cellular Location**

Cytoplasm. Nucleus. Note=Partly colocalized with an endoplasmic reticulum marker, HSP90B1. Shuttles between nucleus and cytoplasm

**Tissue Location**

Isoform 1 is up-regulated in both embryonal rhabdomyosarcoma and alveolar rhabdomyosarcoma cell lines


**FAM193B Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **FAM193B Polyclonal Antibody - Images**



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (FAM193B) Polyclonal Antibody, Unconjugated (bs-14772R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.