

C20orf27 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP55803**Specification**

C20orf27 Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	O9GZN8
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	19291

C20orf27 Polyclonal Antibody - Additional Information**Gene ID** 54976**Other Names**

UPF0687 protein C20orf27, C20orf27

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

C20orf27 Polyclonal Antibody - Protein Information**Name** ADISSP ([HGNC:15873](#))**Function**

Adipocyte-secreted protein (adipokine) that acts as a key regulator for white adipose tissue (WAT) thermogenesis and glucose homeostasis at least in part through activation of protein kinase A (PKA).

Cellular Location

Secreted {ECO:0000250|UniProtKB:Q9D1K7}.


C20orf27 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

C20orf27 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (human stomach); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (C20orf27) Polyclonal Antibody, Unconjugated (bs-15108R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat stomach); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (C20orf27) Polyclonal Antibody, Unconjugated (bs-15108R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.