

**C20orf94 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP55811****Specification**

---

**C20orf94 Polyclonal Antibody - Product Information**

Application	IHC-P
Primary Accession	<a href="#">Q5VYV7</a>
Reactivity	Rat, Pig
Host	Rabbit
Clonality	Polyclonal
Calculated MW	45552

**C20orf94 Polyclonal Antibody - Additional Information****Gene ID** 128710**Other Names**

Protein SLX4IP, SLX4-interacting protein, SLX4IP, C20orf94

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**


Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**C20orf94 Polyclonal Antibody - Protein Information****Name** SLX4IP**Synonyms** C20orf94**C20orf94 Polyclonal Antibody - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**C20orf94 Polyclonal Antibody - Images**



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (C20orf94) Polyclonal Antibody, Unconjugated (bs-15119R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Chromosome 20 open reading frame 94; C20orf94) Polyclonal Antibody, Unconjugated (bs-15119R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.