

C21orf7 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP55816

Specification

C21orf7 Polyclonal Antibody - Product Information

Application WB, IHC-P, IHC-F, IF, ICC, E

Primary Accession P57077

Reactivity Rat, Pig, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 16372

C21orf7 Polyclonal Antibody - Additional Information

Gene ID 56911

Other Names

MAP3K7 C-terminal-like protein, TAK1-like protein, MAP3K7CL, C21orf7, TAK1L

Dilution

 $< span class = "dilution_WB">WB~~1:1000 < /span> < br \> < span class = "dilution_IHC-P">IHC-P~~N/A < /span> < br \> < span class = "dilution_IHC-F">IHC-F~~N/A < /span> < br \> < span class = "dilution_IF">IF~~1:50~200 < /span> < br \> < span class = "dilution_ICC">ICC~~N/A < /span> < br \> < span class = "dilution_E">E~~N/A < /span> < br \> < span class = "dilution_E">E~~N/A < /span> < br \> < span class = "dilution_E">E~~N/A < /span> < br \> < span class = "dilution_E">E~~N/A < /span> < br \> < span class = "dilution_E">E~~N/A < /span> < br \> < span class = "dilution_E">E~~N/A < /span> < br \> < span class = "dilution_E">E~~N/A < /span> < span class = "dilution_E">E~~N/A < /span < span class = "dilution_E">E~~N/A < span < span class =$

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

C21orf7 Polyclonal Antibody - Protein Information

Name MAP3K7CL

Synonyms C21orf7, TAK1L

Tissue Location

Detected in lung and peripheral blood leukocytes. Expressed predominantly in peripheral blood leukocytes and ubiquitously in adult and fetal tissues. Also expressed strongly in breast carcinoma GI-101, colon adenocarcinoma GI-112, and prostatic adenocarcinoma PC3

C21orf7 Polyclonal Antibody - Protocols






Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

C21orf7 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Rat pancreas); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (C21orf7) Polyclonal Antibody, Unconjugated (bs-15128R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.