

C6orf222 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP55892**Specification**

C6orf222 Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	P0C671
Host	Rabbit
Clonality	Polyclonal
Calculated MW	71930

C6orf222 Polyclonal Antibody - Additional Information**Gene ID** 389384**Other Names**

Protein BNIP5, BNIP5 (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=33769)
target="_blank">HGNC:33769), C6orf222

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage


Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

C6orf222 Polyclonal Antibody - Protein Information**Name** BNIP5 ([HGNC:33769](#))**Synonyms** C6orf222**C6orf222 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

C6orf222 Polyclonal Antibody - Images



Sample:

U937 Cell (Human) Lysate at 30 ug

LOVO Cell (Human) Lysate at 30 ug


A549 Cell (Human) Lysate at 30 ug

Primary: Anti-C6orf222 (bs-15241R) at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 42 kD


Observed band size: 42 kD



Tissue/cell: human liver cancer; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-C6orf222 Polyclonal Antibody, Unconjugated(bs-15241R) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: Human lung carcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-C6orf222 Polyclonal Antibody, Unconjugated(bs-15241R) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining