

# GPCR MRGX2 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP55967

## Specification

# **GPCR MRGX2** Polyclonal Antibody - Product Information

Application Primary Accession Host Clonality Calculated MW IHC-P, IHC-F, IF, ICC, E <u>096LB1</u> Rabbit Polyclonal 37099

## GPCR MRGX2 Polyclonal Antibody - Additional Information

Gene ID 117194

**Other Names** Mas-related G-protein coupled receptor member X2, MRGPRX2, MRGX2

Dilution

<span class ="dilution\_IHC-P">IHC-P~~N/A</span><br \><span class ="dilution\_IHC-F">IHC-F~~N/A</span><br \><span class ="dilution\_IF">IF~~1:50~200</span><br \><span class ="dilution\_ICC">ICC~~N/A</span><br \><span class ="dilution\_E">E~~N/A</span>

Format 0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage** Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

## **GPCR MRGX2** Polyclonal Antibody - Protein Information

Name MRGPRX2

Synonyms MRGX2

#### Function

Mast cell-specific receptor for basic secretagogues, i.e. cationic amphiphilic drugs, as well as endoor exogenous peptides, consisting of a basic head group and a hydrophobic core (PubMed:<a href="http://www.uniprot.org/citations/25517090" target="\_blank">25517090</a>). Recognizes and binds small molecules containing a cyclized tetrahydroisoquinoline (THIQ), such as non-steroidal neuromuscular blocking drugs (NMBDs), including tubocurarine and atracurium. In response to these compounds, mediates pseudo-allergic reactions characterized by histamine release, inflammation and airway contraction (By similarity). Acts as a receptor for a number of other ligands, including peptides and alkaloids, such as cortistatin-14, proadrenomedullin N-terminal peptides PAMP-12 and, at lower extent, PAMP-20, antibacterial protein LL-37, PMX-53



peptide, beta-defensins, and complanadine A.

**Cellular Location** Cell membrane; Multi-pass membrane protein

Tissue Location

Mainly expressed in mast cells. Has a limited expression profile, both peripheral and within the central nervous system, with highest levels in dorsal root ganglion (PubMed:12915402) Detected in blood vessels, scattered lymphocytes, and gastrointestinal ganglia (at protein level) (PubMed:16161007)

# **GPCR MRGX2** Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

**GPCR MRGX2 Polyclonal Antibody - Images**