

TRP1 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP56026

Specification

TRP1 Polyclonal Antibody - Product Information

Application WB, IHC-P, IHC-F, IF, ICC, E

Primary Accession
Reactivity
Rat, Bovine
Host
Clonality
Calculated MW
Physical State
P17643
Rat, Bovine
Rabbit
Polyclonal
St KDa
Liquid

Immunogen KLH conjugated synthetic peptide derived

laG

from human TYRP1

Epitope Specificity 101-200/537

Isotype
Purity
affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Melanosome membrane; Single-pass type I

membrane protein (By similarity).
Note=Located to mature stage III and IV
melanosomes and apposed endosomal

tubular membranes. Transported to pigmented melanosomes by the BLOC-1

complex (By similarity).

SIMILARITY

DISEASE

Belongs to the tyrosinase family.

Albinism oculocutaneous 3 (OCA3)

[MIM:203290]: An autosomal recessive disorder in which the biosynthesis of melanin pigment is reduced in skin, hair, and eyes. Tyrosinase activity is normal and patients have only moderate reduction of pigment. The eyes present red reflex on transillumination of the iris, dilution of color of iris, nystagmus and strabismus. Darker-skinned individuals have bright copper-red coloration of the skin and hair. Note=The disease is caused by mutations affecting the gene represented in this

entry.

Important Note This product as supplied is intended for research use only, not for use in human,

therapeutic or diagnostic applications.

Background Descriptions

TRP1 is a melanosomal enzyme that belongs to the tyrosinase family and plays an important role in the melanin biosynthetic pathway. Defects in this gene are the cause of rufous oculocutaneous albinism and oculocutaneous albinism type III.



TRP1 Polyclonal Antibody - Additional Information

Gene ID 7306

Other Names

5, 6-dihydroxyindole-2-carboxylic acid oxidase, DHICA oxidase, 1.14.18.-, Catalase B, Glycoprotein 75, Melanoma antigen gp75, Tyrosinase-related protein 1, TRP, TRP-1, TRP1, TYRP1, CAS2 {ECO:0000303|PubMed:1906272}, TYRP, TYRRP

Target/Specificity

Pigment cells.

Dilution

```
<span class ="dilution_WB">WB~~1:1000</span><br \><span class
="dilution_IHC-P">IHC-P~~N/A</span><br \><span class
="dilution_IHC-F">IHC-F~~N/A</span><br \><span class
="dilution_IF">IF~~1:50~200</span><br \><span class ="dilution_ICC">ICC~~N/A</span><br \><span class ="dilution_ICC">ICC~~N/A</span><br \><span class = "dilution_ICC">ICC~~N/A</span>
```

Storage

Store at -20 $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 $^{\circ}$ C.

TRP1 Polyclonal Antibody - Protein Information

Name TYRP1 (HGNC:12450)

Function

Plays a role in melanin biosynthesis (PubMed:16704458, PubMed:22556244, PubMed:23504663). Catalyzes the oxidation of 5,6- dihydroxyindole-2-carboxylic acid (DHICA) into indole-5,6-quinone-2-carboxylic acid in the presence of bound Cu(2+) ions, but not in the presence of Zn(2+) (PubMed:28661582). May regulate or influence the type of melanin synthesized (PubMed:16704458, PubMed:22556244). Also to a lower extent, capable of hydroxylating tyrosine and producing melanin (By similarity).

Cellular Location

Melanosome membrane {ECO:0000250|UniProtKB:P07147}; Single-pass type I membrane protein {ECO:0000250|UniProtKB:P07147}. Note=Located to mature stage III and IV melanosomes and apposed endosomal tubular membranes. Transported to pigmented melanosomes by the BLOC-1 complex. Proper trafficking to melanosome is regulated by SGSM2, ANKRD27, RAB9A, RAB32 and RAB38 {ECO:0000250|UniProtKB:P07147}

Tissue Location

Pigment cells.


TRP1 Polyclonal Antibody - Protocols



Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

TRP1 Polyclonal Antibody - Images




Sample: B16 (Mouse)cell Lysate at 40 ug

Primary: Anti-TRP1(bs-15510R) at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 58 kD Observed band size: 55 kD



Paraformaldehyde-fixed, paraffin embedded (Human skin); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Tyrosinase-related protein 1) Polyclonal Antibody, Unconjugated (bs-15510R) at 1:400 overnight at 4°C, followed by a conjugated secondary antibody (sp-0023) for 20 minutes and DAB staining.

