

**IFT20 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP56041****Specification**

---

**IFT20 Polyclonal Antibody - Product Information**

Application	FC
Primary Accession	<a href="#">Q8IY31</a>
Reactivity	Rat, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	15281

**IFT20 Polyclonal Antibody - Additional Information****Gene ID** 90410**Other Names**

Intraflagellar transport protein 20 homolog, hIFT20, IFT20

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**IFT20 Polyclonal Antibody - Protein Information****Name** IFT20**Function**

Part of intraflagellar transport (IFT) particles involved in ciliary process assembly (PubMed:<a href="http://www.uniprot.org/citations/17604723" target="\_blank">17604723</a>). May play a role in the trafficking of ciliary membrane proteins from the Golgi complex to the cilium (PubMed:<a href="http://www.uniprot.org/citations/16775004" target="\_blank">16775004</a>). Regulates the platelet-derived growth factor receptor-alpha (PDGFRA) signaling pathway. Required for protein stability of E3 ubiquitin ligases CBL and CBLB that mediate ubiquitination and internalization of PDGFRA for proper feedback inhibition of PDGFRA signaling (PubMed:<a href="http://www.uniprot.org/citations/29237719" target="\_blank">29237719</a>). Essential for male fertility. Plays an important role in spermatogenesis, particularly spermiogenesis, when germ cells form flagella. May play a role in the transport of flagellar proteins ODF2 and SPAG16 to build sperm flagella and in the removal of redundant sperm cytoplasm (By similarity). Also involved in autophagy since it is required for trafficking of ATG16L and the expansion of the autophagic compartment (By similarity).

**Cellular Location**

Golgi apparatus, cis-Golgi network {ECO:0000250|UniProtKB:Q61025}. Cytoplasm, cytoskeleton,

microtubule organizing center, centrosome, centriole {ECO:0000250|UniProtKB:Q61025}. Cytoplasm, cytoskeleton, cilium basal body {ECO:0000250|UniProtKB:Q61025}. Cell projection, cilium {ECO:0000250|UniProtKB:Q61025}. Cytoplasm, cytoskeleton {ECO:0000250|UniProtKB:Q61025}. Golgi apparatus {ECO:0000250|UniProtKB:Q61025}. Cytoplasmic vesicle, secretory vesicle, acrosome {ECO:0000250|UniProtKB:Q61025}. Cytoplasm {ECO:0000250|UniProtKB:Q61025}. Note=Present at the centrosomes during the cell cycle and associated with the proximal portion of the mother centriole and the lateral aspect of the daughter centriole. Associated with basal body at the base of primary cilia. Detected in the Golgi apparatus of round spermatids and late spermatocytes. Also detected in the manchette of step 10-12 spermatids. In step 14 spermatids, found in the basal body of the sperm tail. Localization in the manchette of elongating spermatids is dependent on SPAG17 {ECO:0000250|UniProtKB:Q61025}

### Tissue Location


Expressed in almost all tissues.

### IFT20 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### IFT20 Polyclonal Antibody - Images



Blank control: Mouse spleen.

Primary Antibody (green line): Rabbit Anti-IFT20 antibody (bs-15561R)

Dilution: 2 µg /10<sup>6</sup> cells;

Isotype Control Antibody (orange line): Rabbit IgG .

Secondary Antibody : Goat anti-rabbit IgG-AF488

Dilution: 1 µg /test.

Protocol

The cells were fixed with 70% ethanol (10min at room temperature) and then were incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.