

FAM83G Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP56068**Specification**

FAM83G Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	A6ND36
Reactivity	Rat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	91 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human FAM83G
Epitope Specificity	731-823/823
Isotype	IgG
Purity	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the FAM83 family.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

Chromosome 17 makes up over 2.5% of the human genome with about 81 million bases encoding over 1,200 genes. Two key tumor suppressor genes are associated with chromosome 17, namely, p53 and BRCA1. Tumor suppressor p53 is necessary for maintenance of cellular genetic integrity by moderating cell fate through DNA repair versus cell death. Malfunction or loss of p53 expression is associated with malignant cell growth and Li-Fraumeni syndrome. Like p53, BRCA1 is directly involved in DNA repair, though specifically it is recognized as a genetic determinant of early onset breast cancer and predisposition to cancers of the ovary, colon, prostate gland and fallopian tubes. Chromosome 17 is also linked to neurofibromatosis, a condition characterized by neural and epidermal lesions, and dysregulated Schwann cell growth. Alexander disease, Birt-Hogg-Dube syndrome and Canavan disease are also associated with chromosome 17. The FAM83G gene product has been provisionally designated FAM83G pending further characterization.

FAM83G Polyclonal Antibody - Additional Information**Gene ID** 644815**Other Names**Protein FAM83G, Protein associated with SMAD1, FAM83G, PAWS1
{ECO:0000303|PubMed:24554596}**Dilution**

IHC-P~~N/A<br \><span class
="dilution_IHC-F">IHC-F~~N/A<br \><span class
="dilution_IF">IF~~1:50~200<br \>ICC~~N/A<br
\>E~~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

FAM83G Polyclonal Antibody - Protein Information

Name FAM83G

Synonyms PAWS1 {ECO:0000303|PubMed:24554596}

Function

Substrate for type I BMP receptor kinase involved in regulation of some target genes of the BMP signaling pathway. Also regulates the expression of several non-BMP target genes, suggesting a role in other signaling pathways.

Cellular Location

Cytoplasm, cytosol. Nucleus Note=Detected predominantly in the cytosol. Upon BMP stimulation, a small portion localizes the nucleus.

FAM83G Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FAM83G Polyclonal Antibody - Images