

FBG4 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP56077

Specification

FBG4 Polyclonal Antibody - Product Information

Application
Primary Accession
Reactivity
Host
Clonality
Calculated MW

IHC-P, IHC-F, IF, ICC, E
O96EF6
Rat, Dog, Bovine
Rabbit
Polyclonal
31479

FBG4 Polyclonal Antibody - Additional Information

Gene ID 115290

Other Names

F-box only protein 17, F-box only protein 26, FBXO17, FBG4, FBX17, FBX26, FBXO26

Dilution

IHC-P~~N/A<br \> <span class
="dilution_IHC-F">IHC-F~~N/A<br \> <span class
="dilution_IF">IF~~1:50~200<br \> ICC~~N/A<br \> E~~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 $^{\circ}$ C.

FBG4 Polyclonal Antibody - Protein Information

Name FBXO17


Synonyms FBG4, FBX17, FBX26, FBXO26

Function

Substrate-recognition component of the SCF (SKP1-CUL1-F-box protein)-type E3 ubiquitin ligase complex. Able to recognize and bind denatured glycoproteins, which are modified with complex-type oligosaccharides. Also recognizes sulfated glycans. Does not bind high- mannose glycoproteins.

Tissue Location

Expressed in heart, skeletal muscle, liver and kidney. Expressed at lower levels in spleen and brain





FBG4 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

FBG4 Polyclonal Antibody - Images