

**FGD4 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP56110****Specification**

---

**FGD4 Polyclonal Antibody - Product Information**

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	<a href="#">Q96M96</a>
Reactivity	Rat, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	86626

**FGD4 Polyclonal Antibody - Additional Information****Gene ID** 121512**Other Names**

FYVE, RhoGEF and PH domain-containing protein 4, Actin filament-binding protein frabin, FGD1-related F-actin-binding protein, Zinc finger FYVE domain-containing protein 6, FGD4, FRABP, ZFYVE6

**Dilution**

IHC-P ~ ~ N/A  
IHC-F ~ ~ N/A  
IF ~ ~ 1:50 ~ 200  
ICC ~ ~ N/A  
E ~ ~ N/A

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**FGD4 Polyclonal Antibody - Protein Information****Name** FGD4**Synonyms** FRABP, ZFYVE6**Function**

Activates CDC42, a member of the Ras-like family of Rho- and Rac proteins, by exchanging bound GDP for free GTP. Plays a role in regulating the actin cytoskeleton and cell shape. Activates MAPK8 (By similarity).

**Cellular Location**

Cytoplasm, cytoskeleton. Cell projection, filopodium. Note=Concentrated in filopodia and poorly

detected at lamellipodia. Binds along the sides of actin fibers (By similarity).

**Tissue Location**

Expressed in different tissues, including brain, cerebellum, peripheral nerve, skeletal muscle, heart, uterus, placenta and testis.

**FGD4 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**FGD4 Polyclonal Antibody - Images**