

Focadhesin Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP56159

Specification

Focadhesin Polyclonal Antibody - Product Information

Application IHC-P, IHC-F, IF, ICC

Primary Accession
Reactivity
Rat, Pig, Dog
Host
Clonality
Polyclonal
Calculated MW
200 KDa

Calculated MW 200 KDa
Physical State Liquid

Immunogen KLH conjugated synthetic peptide derived

laG

from human Focadhesin

Epitope Specificity 1-100/1801

Isotype Purity

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Membrane. Cell junction; focal adhesion.

Colocalizes with VCL in astrocytes.

SUBUNIT Interacts with VCL.

Important Note This product as supplied is intended for research use only, not for use in human,

therapeutic or diagnostic applications.

Background Descriptions

affinity purified by Protein A

Chromosome 9 consists of about 145 million bases and 4% of the human genome and encodes nearly 900 genes. Considered to play a role in gender determination, deletion of the distal portion of 9p can lead to development of male to female sex reversal, the phenotype of a female with a male X,Y genotype. Hereditary hemorrhagic telangiectasia, which is characterized by harmful vascular defects, is associated with the chromosome 9 gene encoding endoglin protein, ENG. Familial dysautonomia is also associated with chromosome 9 though through the gene IKBKAP. Notably, chromosome 9 encompasses the largest interferon family gene cluster. Chromosome 9 is partnered with chromosome 22 in the translocation leading to the aberrant production of BCR-ABL fusion protein often found in leukemias. The KIAA1797 gene product has been provisionally designated KIAA1797 pending further characterization.

Focadhesin Polyclonal Antibody - Additional Information

Gene ID 54914

Other Names

Focadhesin, FOCAD, KIAA1797

Target/Specificity

Ubiquitous. High expression in brain followed by testis, muscle, pancreas, heart, ovary, small



intestine, placenta, prostate, thymus, kidney, colon, liver, lung, spleen and leukocytes. Expression is reduced in most glioblastomas and all glioblastoma cell lines.

Dilution

- IHC-P~~N/A<br \><span class</pre>
- ="dilution_IHC-F">IHC-F~~N/A<br \><span class
- ="dilution IF">IF \sim 1:50 \sim 200
span class ="dilution ICC">ICC \sim N/A

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Focadhesin Polyclonal Antibody - Protein Information

Name FOCAD (HGNC:23377)

Synonyms KIAA1797

Function

Required for the maintenance of SKIC2 and SKIC3 proteostatic levels in the liver. May be involved in the regulation of RNA degradation by the exosome complex (PubMed:35864190). Potential tumor suppressor in gliomas.

Cellular Location

Cell junction, focal adhesion. Cytoplasm, cytosol. Note=In astrocytes, colocalizes with VCL to the end of actin stress fibers, which normally terminate at focal adhesions. In hepatocytes, it is found in the cytosol

Tissue Location

Ubiquitous. High expression in brain followed by testis, muscle, pancreas, heart, ovary, small intestine, placenta, prostate, thymus, kidney, colon, liver, lung, spleen and leukocytes Expression is reduced in most glioblastomas and all glioblastoma cell lines.

Focadhesin Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Focadhesin Polyclonal Antibody - Images