

FOXI3 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP56161

Specification

FOXI3 Polyclonal Antibody - Product Information

Application IHC-P, IHC-F, IF, ICC

Primary Accession
Reactivity
Rat, Bovine
Host
Clonality
Calculated MW
Physical State

A8MTJ6
Rat, Bovine
Rabbit
Polyclonal
Liquid

Immunogen KLH conjugated synthetic peptide derived

from human FOXI3

Epitope Specificity 121-220/420

Isotype IgG
Purity

affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Nucleus.

SIMILARITY
Important Note
Contains 1 fork-head DNA-binding domain.
This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

FOXI3 Polyclonal Antibody - Additional Information

Gene ID 344167

Other Names

Forkhead box protein I3, FOXI3

Dilution

IHC-P~~N/A<br \><span class</pre>

="dilution IHC-F">IHC-F~~N/A<br \><span class

="dilution IF">IF~~1:50~200
or \>ICC~~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 $^{\circ}$ C.

FOXI3 Polyclonal Antibody - Protein Information



Name FOXI3 {ECO:0000303|PubMed:36260083, ECO:0000312|HGNC:HGNC:35123}

Function

Transcription factor required for pharyngeal arch development, which is involved in hair, ear, jaw and dental development (PubMed:37041148). May act as a pioneer transcription factor during pharyngeal arch development (By similarity). Required for epithelial cell differentiation within the epidermis (By similarity). Acts at multiple stages of otic placode induction: necessary for preplacodal ectoderm to execute an inner ear program (By similarity). Required for hair follicle stem cell specification (By similarity). Acts downstream of TBX1 for the formation of the thymus and parathyroid glands from the third pharyngeal pouch (By similarity).

Cellular Location Nucleus.

FOXI3 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

FOXI3 Polyclonal Antibody - Images