

**ZNF782 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP56270****Specification**

---

**ZNF782 Polyclonal Antibody - Product Information**

Application	IHC-P
Primary Accession	<a href="#">Q6ZMW2</a>
Host	Rabbit
Clonality	Polyclonal
Calculated MW	80904

**ZNF782 Polyclonal Antibody - Additional Information****Gene ID** 158431**Other Names**

Zinc finger protein 782, ZNF782

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**ZNF782 Polyclonal Antibody - Protein Information****Name** ZNF782**Function**

May be involved in transcriptional regulation.

**Cellular Location**

Nucleus.


**ZNF782 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

- [Flow Cytometry](#)
- [Cell Culture](#)

#### **ZNF782 Polyclonal Antibody - Images**



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ZNF782) Polyclonal Antibody, Unconjugated (bs-16456R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.