

ZNF777 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP56272**Specification**

ZNF777 Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	O9ULD5
Reactivity	Rat, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	93762

ZNF777 Polyclonal Antibody - Additional Information**Gene ID** 27153**Other Names**

Zinc finger protein 777, ZNF777, KIAA1285

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

ZNF777 Polyclonal Antibody - Protein Information**Name** ZNF777 ([HGNC:22213](#))**Synonyms** KIAA1285**Function**

May be involved in transcriptional repression (PubMed:31856708). Inhibits cell proliferation through CDKN1A/p21 induction by down-regulation of NIBAN1/FAM129A at low cell density (PubMed:25560148).

Cellular Location


Nucleus.

ZNF777 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZNF777 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (human brain glioma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ZNF777) Polyclonal Antibody, Unconjugated (bs-16460R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.