

Hemoglobin subunit delta Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP56276**Specification**

Hemoglobin subunit delta Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	P02042
Host	Rabbit
Clonality	Polyclonal
Calculated MW	16055

Hemoglobin subunit delta Polyclonal Antibody - Additional Information**Gene ID** 3045**Other Names**

Hemoglobin subunit delta, Delta-globin, Hemoglobin delta chain, HBD

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Hemoglobin subunit delta Polyclonal Antibody - Protein Information**Name** HBD**Function**

Involved in oxygen transport from the lung to the various peripheral tissues.

Tissue Location

Red blood cells.


Hemoglobin subunit delta Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

- [Flow Cytometry](#)
- [Cell Culture](#)

Hemoglobin subunit delta Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Human kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (HBD) Polyclonal Antibody, Unconjugated (bs-16467R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.