

ZNF770 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP56283**Specification**

ZNF770 Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q6IQ21
Reactivity	Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	80007

ZNF770 Polyclonal Antibody - Additional Information**Gene ID** 54989**Other Names**

Zinc finger protein 770, ZNF770

Dilution

IHC-P ~ ~ N/A
IHC-F ~ ~ N/A
IF ~ ~ 1:50 ~ 200
ICC ~ ~ N/A
E ~ ~ N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

ZNF770 Polyclonal Antibody - Protein Information**Name** ZNF770**Function**

May be involved in transcriptional regulation.

Cellular Location


Nucleus.

ZNF770 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZNF770 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ZNF770) Polyclonal Antibody, Unconjugated (bs-16483R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.