

HSF4 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP56314

Specification

HSF4 Polyclonal Antibody - Product Information

Application Primary Accession Reactivity Host Clonality Calculated MW WB, IHC-P, IHC-F, IF, ICC, E <u>O9ULV5</u> Rat, Pig, Dog, Bovine Rabbit Polyclonal 53011

HSF4 Polyclonal Antibody - Additional Information

Gene ID 3299

Other Names Heat shock factor protein 4, HSF 4, hHSF4, Heat shock transcription factor 4, HSTF 4, HSF4

Dilution WB~~1:1000<br \>IHC-P~~N/A<br \>IHC-F~~N/A<br \>IF~~1:50~200<br \>ICC~~N/A<br \>ICC~~N/A<br \>ICC~~N/A

Format 0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

HSF4 Polyclonal Antibody - Protein Information

Name HSF4

Function

Heat-shock transcription factor that specifically binds heat shock promoter elements (HSE) (PubMed:22587838, PubMed:23507146). Required for denucleation and organelle rupture and degradation that occur during eye lens terminal differentiation, when fiber cells that compose the lens degrade all membrane-bound organelles in order to provide lens with transparency to allow the passage of light (By similarity). In this process, may regulate denucleation of lens fiber cells in part by activating DNASE2B transcription (By similarity). May be involved in DNA repair through the transcriptional regulation of RAD51 (PubMed:22587838" target="_blank">22587838" target="_blank">22587838" target="_blank">22587838" target="_blank">22587838



target="_blank">22587838). May up-regulate p53/TP53 protein in eye lens fiber cells, possibly through protein stabilization (PubMed:28981088). In the eye lens, controls the expression of alpha-crystallin B chain/CRYAB and consequently may be involved in the regulation of lysosomal acidification (By similarity).

Cellular Location Nucleus.

Tissue Location Expressed in heart, skeletal muscle, eye and brain, and at much lower levels in some other tissues

HSF4 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

HSF4 Polyclonal Antibody - Images