

**HSF4 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP56314****Specification**

---

**HSF4 Polyclonal Antibody - Product Information**

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	<a href="#">O9ULV5</a>
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	53 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human HSF4
Epitope Specificity	21-120/492
Isotype	IgG
<b>Purity</b>	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nucleus.
SIMILARITY	Belongs to the HSF family.
Post-translational modifications	Phosphorylated mainly on serine residues. Phosphorylation on Ser-298 promotes sumoylation on Lys-293. Isoform HSF4B is constitutively sumoylated. Sumoylation represses the transcriptional activity and is promoted by phosphorylation on Ser-298. HSFA is not sumoylated.
DISEASE	Defects in HSF4 are the cause of cataract zonular HSF4-related (CZ-HSF4) [MIM:116800]. A form of zonular cataract. Zonular or lamellar cataracts are opacities, broad or narrow, usually consisting of powdery white dots affecting only certain layers or zones between the cortex and nucleus of an otherwise clear lens. The opacity may be so dense as to render the entire central region of the lens completely opaque, or so translucent that vision is hardly if at all impeded. Zonular cataracts generally do not involve the embryonic nucleus, though sometimes they involve the fetal nucleus. Usually sharply separated from a clear cortex outside them, they may have projections from their outer edges known as riders or spokes. Defects in HSF4 are the cause of cataract Marner type (CAM) [MIM:116800].

**A form of cataract with variable and progressive opacities. Affected individuals present with zonular cataract, although some have nuclear, anterior polar, or stellate cataract. Finger malformation is observed in some kindreds.**

#### Important Note

**This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.**

#### Background Descriptions

Heat-shock transcription factors (HSFs) activate heat-shock response genes under conditions of heat or other stresses. HSF4 lacks the carboxyl-terminal hydrophobic repeat which is shared among all vertebrate HSFs and has been suggested to be involved in the negative regulation of DNA binding activity. Two alternatively spliced transcripts encoding distinct isoforms and possessing different transcriptional activity have been described. [provided by RefSeq, Jul 2008]

### HSF4 Polyclonal Antibody - Additional Information

**Gene ID** 3299

#### Other Names

Heat shock factor protein 4, HSF 4, hHSF4, Heat shock transcription factor 4, HSTF 4, HSF4

#### Target/Specificity

Expressed in heart, skeletal muscle, eye and brain, and at much lower levels in some other tissues.

#### Dilution

WB~1:1000  
IHC-P~N/A  
IHC-F~N/A  
IF~1:50~200  
ICC~N/A  
E~N/A

#### Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

#### Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

### HSF4 Polyclonal Antibody - Protein Information

**Name** HSF4

#### Function

Heat-shock transcription factor that specifically binds heat shock promoter elements (HSE) (PubMed: [22587838](http://www.uniprot.org/citations/22587838)), PubMed: [23507146](http://www.uniprot.org/citations/23507146)). Required for denucleation and organelle rupture and degradation that occur during eye lens terminal differentiation, when fiber cells that compose the lens degrade all membrane-bound organelles in order to provide lens with transparency to allow the passage of light (By similarity). In this process, may regulate denucleation of lens fiber cells in part by activating DNASE2B

transcription (By similarity). May be involved in DNA repair through the transcriptional regulation of RAD51 (PubMed:<a href="http://www.uniprot.org/citations/22587838" target="\_blank">22587838</a>). May up-regulate p53/TP53 protein in eye lens fiber cells, possibly through protein stabilization (PubMed:<a href="http://www.uniprot.org/citations/28981088" target="\_blank">28981088</a>). In the eye lens, controls the expression of alpha-crystallin B chain/CRYAB and consequently may be involved in the regulation of lysosomal acidification (By similarity).

**Cellular Location**

Nucleus.

**Tissue Location**

Expressed in heart, skeletal muscle, eye and brain, and at much lower levels in some other tissues

**HSF4 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**HSF4 Polyclonal Antibody - Images**