

TRAPPC5 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP56325**Specification**

TRAPPC5 Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	Q8IUQ0
Reactivity	Rat, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	20783

TRAPPC5 Polyclonal Antibody - Additional Information**Gene ID** 126003**Other Names**

Trafficking protein particle complex subunit 5, TRAPPC5

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

TRAPPC5 Polyclonal Antibody - Protein Information**Name** TRAPPC5**Function**

May play a role in vesicular transport from endoplasmic reticulum to Golgi.

Cellular Location

Golgi apparatus, cis-Golgi network. Endoplasmic reticulum


TRAPPC5 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)

- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TRAPPC5 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (rat heart tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (TRAPPC5) Polyclonal Antibody, Unconjugated (bs-16586R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.