

## **TNFAIP8L1 Polyclonal Antibody**

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP56331

## **Specification**

# **TNFAIP8L1 Polyclonal Antibody - Product Information**

Application IHC-P, IHC-F, IF, ICC, E

Primary Accession <u>Q8WVP5</u>

Reactivity Rat, Dog, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 20827

# TNFAIP8L1 Polyclonal Antibody - Additional Information

#### Gene ID 126282

#### **Other Names**

Tumor necrosis factor alpha-induced protein 8-like protein 1, TIPE1, TNF alpha-induced protein 8-like protein 1, TNFAIP8-like protein 1, Oxidative stress-regulated gene-beta, Oxy-beta, TNFAIP8L1

## **Dilution**

<span class ="dilution\_IHC-P">IHC-P~~N/A</span><br \><span class
="dilution\_IHC-F">IHC-F~~N/A</span><br \><span class
="dilution\_IF">IF~~1:50~200</span><br \><span class ="dilution\_ICC">ICC~~N/A</span><br \><span class ="dilution\_E">E~~N/A</span>

#### **Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

### **Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

# **TNFAIP8L1 Polyclonal Antibody - Protein Information**

# Name TNFAIP8L1

## **Function**


Acts as a negative regulator of mTOR activity.

#### **Cellular Location**

Cytoplasm {ECO:0000250|UniProtKB:Q8K288}.

## **Tissue Location**

High expression detected in most carcinoma cell lines, especially in cells transformed with virus genomes





**TNFAIP8L1** Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

**TNFAIP8L1 Polyclonal Antibody - Images**