

**KPNA3 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP56408****Specification****KPNA3 Polyclonal Antibody - Product Information**

Application	WB, IHC-P, IHC-F, IF, ICC
Primary Accession	<a href="#">000505</a>
Host	Rabbit
Clonality	Polyclonal
Calculated MW	57811

**KPNA3 Polyclonal Antibody - Additional Information****Gene ID** 3839**Other Names**

Importin subunit alpha-4, Importin alpha Q2, Qip2, Karyopherin subunit alpha-3, SRP1-gamma, KPNA3, QIP2

**Dilution**

<span class = "dilution\_WB">WB~~1:1000</span><br /><span class = "dilution\_IHC-P">IHC-P~~N/A</span><br /><span class = "dilution\_IHC-F">IHC-F~~N/A</span><br /><span class = "dilution\_IF">IF~~1:50~200</span><br /><span class = "dilution\_ICC">ICC~~N/A</span>

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**KPNA3 Polyclonal Antibody - Protein Information****Name** KPNA3**Synonyms** QIP2**Function**

Functions in nuclear protein import as an adapter protein for nuclear receptor KPNB1. Binds specifically and directly to substrates containing either a simple or bipartite NLS motif. Docking of the importin/substrate complex to the nuclear pore complex (NPC) is mediated by KPNB1 through binding to nucleoporin FxFG repeats and the complex is subsequently translocated through the pore by an energy requiring, Ran-dependent mechanism. At the nucleoplasmic side of the NPC, Ran binds to importin-beta and the three components separate and importin-alpha and -beta are re-exported from the nucleus to the cytoplasm where GTP hydrolysis releases Ran from importin. The directionality of nuclear import is thought to be conferred by an asymmetric distribution of the GTP- and GDP-bound forms of Ran between the cytoplasm and nucleus. In vitro, mediates the nuclear import of human cytomegalovirus UL84 by recognizing a non-classical NLS. Recognizes NLSs of influenza A virus nucleoprotein probably through ARM repeats 7-9.

**Cellular Location**

Cytoplasm. Nucleus

**Tissue Location**

Ubiquitous. Highest levels in heart and skeletal muscle

**KPNA3 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**KPNA3 Polyclonal Antibody - Images**