

KRT74 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP56427**Specification**

KRT74 Polyclonal Antibody - Product Information

| | |
|-------------------|------------------------|
| Application | IHC-P, WB |
| Primary Accession | Q7RTS7 |
| Reactivity | Dog |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 57865 |

KRT74 Polyclonal Antibody - Additional Information**Gene ID** 121391**Other Names**

Keratin, type II cytoskeletal 74, Cytokeratin-74, CK-74, Keratin-5c, K5C, Keratin-74, K74, Type II inner root sheath-specific keratin-K6irs4, Type-II keratin Kb37, KRT74, K6IRS4, KB37, KRT5C, KRT6IRS4

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

KRT74 Polyclonal Antibody - Protein Information**Name** KRT74**Synonyms** K6IRS4, KB37, KRT5C, KRT6IRS4**Function**

Has a role in hair formation. Specific component of keratin intermediate filaments in the inner root sheath (IRS) of the hair follicle (Probable).

Tissue Location


Highly expressed in hair follicles from scalp. In hair, it is specifically present in the inner root sheath (IRS) of the hair follicle. Present in the IRS Huxley layer, but not in Henle layer or cuticle of the IRS. In the IRS Huxley layer, it is expressed in specialized Huxley cells, termed 'Fluegelzellen, along the area of differentiated Henle cells (at protein level)

KRT74 Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

KRT74 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (KRT74) Polyclonal Antibody, Unconjugated (bs-16833R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Sample: A431 Cell (Human) Lysate at 40 ug
Primary: Anti-KRT74 (bs-16833R) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 58 kD
Observed band size: 55 kD