

KTEL1 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP56443**Specification****KTEL1 Polyclonal Antibody - Product Information**

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q8NBL1
Reactivity	Rat, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	43 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human KTEL1
Epitope Specificity	201-300/392
Isotype	IgG
Purity	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Endoplasmic reticulum lumen.
SIMILARITY	Belongs to the glycosyltransferase 90 family.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

CLP46 is a KTEL motif-containing protein that belongs to the CAP10 family. The KTEL motif at the C-terminus of CLP46 is an endoplasmic reticulum (ER) retention signal which localizes the CLP46 protein to the lumen of the ER. CLP46 is thought to promote cell proliferation and is also believed to be involved with hepatic functions. The Drosophila protein rumi, a CAP10 protein and likely similar in function to CLP46, uses glucosylation to act as a Notch signaling regulator. Expressed in varying degrees in most adult tissue, CLP46 is especially abundant in the liver. CLP46 is not expressed in detectable levels in colon, thymus or small intestine. Human CLP46 shares 94% and 91% sequence similarity with its bovine and mouse homologs, respectively.

KTEL1 Polyclonal Antibody - Additional Information**Gene ID** 56983**Other Names**

Protein O-glucosyltransferase 1, 2.4.1.376, CAP10-like 46 kDa protein, hCLP46, KTEL motif-containing protein 1, Myelodysplastic syndromes relative protein, O-glucosyltransferase Rumi homolog, hRumi, Protein O-xylosyltransferase POGLUT1, 2.4.2.63, POGLUT1 (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=22954)
HGNC:22954

Target/Specificity

Expressed in most adult tissues at different intensities. Abundantly expressed in liver. Expressed also in brain, heart, skeletal muscle, spleen, kidney, placenta, lung and peripheral blood leukocyte. Not detectable in colon, thymus and small intestine.

Dilution

WB~~1:1000<br \>IHC-P~~N/A<br \>IHC-F~~N/A<br \>IF~~1:50~200<br \>ICC~~N/A<br \>E~~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glycerol

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

KTEL1 Polyclonal Antibody - Protein Information

Name POGLUT1 ([HGNC:22954](#))

Function

Dual specificity glycosyltransferase that catalyzes the transfer of glucose and xylose from UDP-glucose and UDP-xylose, respectively, to a serine residue found in the consensus sequence of C- X-S-X-P-C (PubMed:21081508, PubMed:21490058, PubMed:21949356, PubMed:27807076, PubMed:28775322). Specifically targets extracellular EGF repeats of protein such as CRB2, F7, F9 and NOTCH2 (PubMed:21081508, PubMed:21490058, PubMed:21949356, PubMed:27807076, PubMed:28775322). Acts as a positive regulator of Notch signaling by mediating O-glucosylation of Notch, leading to regulate muscle development (PubMed:27807076). Notch glucosylation does not affect Notch ligand binding (PubMed:21490058). Required during early development to promote gastrulation: acts by mediating O-glucosylation of CRB2, which is required for CRB2 localization to the cell membrane (By similarity).

Cellular Location

Endoplasmic reticulum lumen

Tissue Location


Expressed in most adult tissues at different intensities. Abundantly expressed in liver. Expressed also in brain, heart, skeletal muscle, spleen, kidney, placenta, lung and peripheral blood leukocyte. Not detectable in colon, thymus and small intestine Expressed in the epidermis, especially in the upper parts, stratum spinosum and stratum granulosum (at protein level)

KTEL1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

KTEL1 Polyclonal Antibody - Images



Sample: Cerebrum (Mouse) Lysate at 40 ug
Primary: Anti-KTEL1 (bs-16861R) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 43 kD
Observed band size: 43 kD