

HSP20 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) **Catalog # AP56455**

Specification

HSP20 Polyclonal Antibody - Product Information

Application **Primary Accession**

Reactivity Host Clonality Calculated MW **Physical State** Immunogen

Epitope Specificity

Isotype **Purity**

affinity purified by Protein A

6-80/160

KLH conjugated synthetic peptide derived

laG

014558

Rabbit

17 KDa

Liquid

Polyclonal

IHC-P, IHC-F, IF, E

Rat, Dog, Bovine

from human HSPB6

Buffer

SUBCELLULAR LOCATION

SIMILARITY

SUBUNIT

Post-translational modifications

Important Note

0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

Cytoplasm. Nucleus. Note=Translocates to

nuclear foci during heat shock.

Belongs to the small heat shock protein

(HSP20) family.

Homodimer (By similarity). The N-terminus is blocked.

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

Hsp20 is a small heat shock protein related to Hsp25, Hsp27 and may form different hetercomplexes with these proteins. The specific physiological function of Hsp20 is not yet known. It is distributed ubiquitously in tissues, but is found in higher levels in skeletal, smooth and heart muscle. Under normal conditions, Hsp20 is diffusely distributed in the cytosol, but under heat stress conditions, it translocates to the nucleus. Unlike other heat shock proteins the amount of Hsp20 does not increase after heat shock. The Hsp20 was demonstrated to constitute up to 1.3% of the total cellular protein in vertebrate tissues, especially in muscle, and its expression is related to muscle contraction, specifically in slow-twitch muscle. Hsp20 may form different heterocomplexes with other Hsp's, such as alpha-crystalline and Hsp25. Phosphorylated form of Hsp20 is proposed to interact with monomeric actin whereas dephosphorylated form binds polymeric actin filaments. In normal conditions Hsp20 is diffusely disturbed in cytosol but under the heat stress it undergoes translocation to membrane fraction.

HSP20 Polyclonal Antibody - Additional Information

Gene ID 126393



Other Names

Heat shock protein beta-6, HspB6, Heat shock 20 kDa-like protein p20, HSPB6

Dilution

- IHC-P~~N/A<br \>IHC-F~~N/A<br \><span class
- ="dilution IF">IF \sim 1:50 \sim 200
span class ="dilution E">E \sim N/A

Storage

Store at -20 $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 $^{\circ}$ C.

HSP20 Polyclonal Antibody - Protein Information

Name HSPB6

Function

Small heat shock protein which functions as a molecular chaperone probably maintaining denatured proteins in a folding- competent state. Seems to have versatile functions in various biological processes. Plays a role in regulating muscle function such as smooth muscle vasorelaxation and cardiac myocyte contractility. May regulate myocardial angiogenesis implicating KDR. Overexpression mediates cardioprotection and angiogenesis after induced damage. Stabilizes monomeric YWHAZ thereby supporting YWHAZ chaperone-like activity.

Cellular Location


Cytoplasm. Nucleus. Secreted Note=Translocates to nuclear foci during heat shock

HSP20 Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

HSP20 Polyclonal Antibody - Images







Paraformaldehyde-fixed, paraffin embedded (human breast); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (HSP20) Polyclonal Antibody, Unconjugated (bs-1688R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.