

KCTD19 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP56469**Specification**

KCTD19 Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	Q17RG1
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	104938

KCTD19 Polyclonal Antibody - Additional Information**Gene ID** 146212**Other Names**

BTB/POZ domain-containing protein KCTD19, KCTD19

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

KCTD19 Polyclonal Antibody - Protein Information**Name** KCTD19**Function**

Transcription regulator which is essential for male fertility and for the completion of meiotic prophase in spermatocytes. Regulates progression of the pachytene stage of meiotic prophase and promotes the transcriptional activation activity ZNF541. Required for the organization of chromosomes during metaphase I.

Cellular Location

Nucleus {ECO:0000250|UniProtKB:Q562E2}.


KCTD19 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


KCTD19 Polyclonal Antibody - Images



Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-KCTD19 Polyclonal Antibody, Unconjugated(bs-16930R) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: mouse testis tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-KCTD19 Polyclonal Antibody, Unconjugated(bs-16930R) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining