

KDM5A Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP56474**Specification**

KDM5A Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	P29375
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	192095

KDM5A Polyclonal Antibody - Additional Information**Gene ID** 5927**Other Names**

Lysine-specific demethylase 5A, 1.14.11.67, Histone demethylase JARID1A, Jumonji/ARID domain-containing protein 1A, Retinoblastoma-binding protein 2, RBBP-2, [histone H3]-trimethyl-L-lysine(4) demethylase 5A, KDM5A (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=9886 target="_blank">HGNC:9886)

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

KDM5A Polyclonal Antibody - Protein Information**Name** KDM5A ([HGNC:9886](#))**Function**

Histone demethylase that specifically demethylates 'Lys-4' of histone H3, thereby playing a central role in histone code. Does not demethylate histone H3 'Lys-9', H3 'Lys-27', H3 'Lys-36', H3 'Lys-79' or H4 'Lys-20'. Demethylates trimethylated and dimethylated but not monomethylated H3 'Lys-4'. Regulates specific gene transcription through DNA-binding on 5'-CCGCCC-3' motif (PubMed:18270511). May stimulate transcription mediated by nuclear receptors. Involved in transcriptional regulation of Hox proteins during cell differentiation (PubMed:19430464). May participate in transcriptional repression of cytokines such as CXCL12. Plays a role in the regulation of the circadian rhythm and in maintaining the normal periodicity of the circadian clock. In a histone demethylase-independent manner, acts as a coactivator of the CLOCK-BMAL1-mediated transcriptional activation of PER1/2 and other clock-controlled genes and increases histone acetylation at PER1/2 promoters by inhibiting the

activity of HDAC1 (By similarity). Seems to act as a transcriptional corepressor for some genes such as MT1F and to favor the proliferation of cancer cells (PubMed:27427228).

Cellular Location


Nucleus, nucleolus. Nucleus {ECO:0000250|UniProtKB:Q3UXZ9} Note=Occupies promoters of genes involved in RNA metabolism and mitochondrial function. {ECO:0000250|UniProtKB:Q3UXZ9}

KDM5A Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

KDM5A Polyclonal Antibody - Images



Tissue/cell: human laryngocarcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded;
Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;
Incubation: Anti-KDM5A Polyclonal Antibody, Unconjugated(bs-16947R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining