

### KIF6 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP56538

# **Specification**

### KIF6 Polyclonal Antibody - Product Information

Application IHC-P, IHC-F, IF, ICC, E

Primary Accession
Reactivity
Rat, Pig
Host
Clonality
Calculated MW
Rat, Pig
Rabbit
Polyclonal
92569

### KIF6 Polyclonal Antibody - Additional Information

**Gene ID 221458** 

#### **Other Names**

Kinesin-like protein KIF6, KIF6, C6orf102

#### **Dilution**

<span class ="dilution\_IHC-P">IHC-P~~N/A</span><br \> <span class
="dilution\_IHC-F">IHC-F~~N/A</span><br \> <span class
="dilution\_IF">IF~~1:50~200</span><br \> <span class ="dilution\_ICC">ICC~~N/A</span><br \> <span class ="dilution\_E">E~~N/A</span>

#### **Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

#### **Storage**

Store at -20  $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4  $^{\circ}$ C.

# KIF6 Polyclonal Antibody - Protein Information

Name KIF6

Synonyms C6orf102

### **Cellular Location**

Cytoplasm, cytoskeleton.


## **KIF6 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.




- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

### KIF6 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (mouse brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (KIF6) Polyclonal Antibody, Unconjugated (bs-17057R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Human kidney tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (KIF6) Polyclonal Antibody, Unconjugated (bs-17057R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.