

**KAZALD1 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP56550****Specification****KAZALD1 Polyclonal Antibody - Product Information**

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	<a href="#">Q96I82</a>
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	30 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human KAZALD1
Epitope Specificity	71-170/304
Isotype	IgG
<b>Purity</b>	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Secreted > extracellular space > extracellular matrix.
SIMILARITY	Contains 1 Ig-like C2-type (immunoglobulin-like) domain. Contains 1 IGFBP N-terminal domain. Contains 1 Kazal-like domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

**Background Descriptions**

BONO1, also known as KAZALD1 (kazal-type serine protease inhibitor domain-containing protein 1), IGFBP-rP10, FKSG28 or FKSG40, is a 304 amino acid secreted extracellular matrix protein that promotes matrix assembly. BONO1 is expressed in developing bones and odontoblasts in teeth, where it plays a role in osteoblast proliferation during bone formation and regeneration. BONO1 is also expressed at high levels in spleen, and is found at lower levels in lung, skin, urinary bladder, brain, tongue, kidney and large intestine. Existing as two alternatively spliced isoforms, BONO1 contains one Kazal-like domain, an IGFBP N-terminal domain and a single Ig-like C2-type (immunoglobulin-like) domain. The gene encoding BONO1 maps to human chromosome 10, which contains over 800 genes and 135 million nucleotides. Cockayne syndrome, Cockayne syndrome and trisomy 10 are associated with defects in chromosome 10.

**KAZALD1 Polyclonal Antibody - Additional Information****Gene ID** 81621**Other Names**

Kazal-type serine protease inhibitor domain-containing protein 1, KAZALD1

**Dilution**

<span class = "dilution\_WB">WB~~1:1000</span><br \><span class = "dilution\_IHC-P">IHC-P~~N/A</span><br \><span class = "dilution\_IHC-F">IHC-F~~N/A</span><br \><span class = "dilution\_IF">IF~~1:50~200</span><br \><span class = "dilution\_ICC">ICC~~N/A</span><br \><span class = "dilution\_E">E~~N/A</span>

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**KAZALD1 Polyclonal Antibody - Protein Information**

**Name** KAZALD1

**Function**

Involved in the proliferation of osteoblasts during bone formation and bone regeneration. Promotes matrix assembly (By similarity).


**Cellular Location**


Secreted, extracellular space, extracellular matrix

**KAZALD1 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**KAZALD1 Polyclonal Antibody - Images**



**Sample:**


DU145(Human) Cell Lysate at 40 ug

Primary: Anti-KAZALD1 (bs-17082R) at 1/300 dilution


Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 30 kD

Observed band size: 30 kD



Paraformaldehyde-fixed, paraffin embedded (Human brain glioma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (KAZALD1) Polyclonal Antibody, Unconjugated (bs-17082R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rat small intestine); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (KAZALD1) Polyclonal Antibody, Unconjugated (bs-17082R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.