

TRIM52 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP56565**Specification**

TRIM52 Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	O96A61
Host	Rabbit
Clonality	Polyclonal
Calculated MW	34653

TRIM52 Polyclonal Antibody - Additional Information**Gene ID** 84851**Other Names**

E3 ubiquitin-protein ligase TRIM52, 2.3.2.27, RING finger protein 102, Tripartite motif-containing protein 52, TRIM52, RNF102

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

TRIM52 Polyclonal Antibody - Protein Information**Name** TRIM52**Synonyms** RNF102**Function**

E3 ubiquitin-protein ligase (PubMed:27667714). Positively regulates the NF-kappa-B signaling pathway (PubMed:27667714, PubMed:28073078).

Cellular Location


Cytoplasm. Cytoplasm, cytosol. Nucleus

TRIM52 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TRIM52 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (TRIM52) Polyclonal Antibody, Unconjugated (bs-17115R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.